

Fixed Points of Quasi Contractive Mapping using Ishikawa Iteration

Kasturi Mishra¹, Hadibandhu Pattnaik²

¹Research Scholar, ²Ex-reader,
Dept. of Mathematics, Ravenshaw University, Cuttack, Odisha, India 753003.

Corresponding Author: Kasturi Mishra

Received: 11/07/2016

Revised: 17/07/2016

Accepted: 25/07/2016

ABSTRACT

A map has a fixed point at P. If fixed point theorems have useful applications in analysis. Some of the iterative methods which have been studied are related to S. Banach, W.R. Mann, J. Riemermann, W.G. Daston and a host of other mathematicians.

Studies by Prof. S. Ishikawa and Prof. B.E. Rhoads, throw new light on the iteration process of W.R. Mann, Prof. Ishikawa studied by the following iteration process.

For a subset E of an Ailbert space H, if and only if the sequence generated by $x_{n-1} = (1 - c_n)x_n + c_n T x_n, n \geq 1$. Where (c_n) are real sequence in $[0, 1]$.

Key Words: Fixed point, metric space, picard iteration, Ishikawa iteration.

1.1. INTRODUCTION

Prof B.E. Khoades has shown in [17] that from amongst various generalizations of Banach's contraction principle the definition of quasi contraction by circle [1] is one of the most general contractive definition for which Picard's Iteration gives a unique fixed point. We recall the definition of a Quasi contractive method which states that if there exists a constant $k, 0 \leq k < 1$ such that for each $x, y \in E. \| T_x - T_y \| \leq k \max \{ \| x - y \|, \| x - T_x \|, \| y - T_y \|, \| y - T_x \| \} \dots \dots (x)$. In [3] Hu." has shown that most of the results of [17] which use $M(x_{11}, C_n, T)$ can be extended to Ishikawa's iteration scheme $I(X_1, C_n, d_n, T)$. Thus $I(X_1, C_n, d_n, T)$ becomes a larger class of fixed point iteration method. However he [4 theorem 9] posed an open question whether mann iterative process $M(X_1, C_n, T)$ can be

replaced by that of Ishikawa $I(X_1, C_n, d_n, T)$ for quasi contractive mapping in this chapter our purpose is to show that in a Hilbert space $I(X_1, C_n, d_n, T)$ converges to the fixed point of a Quasi contractive map. This is embodied is theorem I below. Theorem 2 provides a generalization of theorem.

1.2 MAIN RESULTS

Theorem: Let E be a compact and convex subset of a Hilbert space. H and T be a quasi contractive self map on E. Let a sequence (X_n) be defined iteratively on E by

$$X_1 \in E, X_{n+1} = (1 - C_n) X_n + C_n T [(1 - d_n) X_n + d_n T X_n] \dots \dots (1)$$

Where (C_n) and (T_n) are sequences of real numbers such that

(i) $0 \leq C_n \leq d_n \leq 1$

(ii) $\lim_{n \rightarrow \infty} d_n = 0$

$$(iii) \quad \sum_{n=1}^{\infty} C_n d_n = \infty$$

Then (X_n) converges to the fixed point of T.

Pf : Since T is quasi contractive by circle [1] it has a unique fixed point P say. Hence from (*) putting P for y we have for each $x \in E$,

$$\|T_x - P\| \leq k \max \{ \|x - p\|, \|x - T_x\| \} \quad (2)$$

$$\text{writing } Y_n = (1-d_n)X_n + d_n T_{yn} \quad (3)$$

we can express X_{n+1} in (1) as

$$X_{n+1} = (1 - C_n) X_n + C_n T_{yn} \quad (4)$$

We know [33] that for any X, Y, Z in a Hilbert space and for any real number t.

$$\|t_x + (1-t)y - z\|^2 = t \|x - z\|^2 + (1-t) \|y - z\|^2 - t(1-t) \|x - y\|^2 \quad (5)$$

Hence from (3) and (4) we have the following relation.

$$\|x_{n+1} - p\|^2 = 1 - C_n \|X_n - p\|^2 + C_n \|T_{yn} - p\|^2 - C_n (1 - C_n) \|X_n - T_{yn}\|^2 \quad (6)$$

$$\|Y_n - T_{yn}\|^2 = (1-d_n) \|X_n - T_{yn}\|^2 + d_n \|T_{xn} - T_{yn}\|^2 - d_n (1-d_n) \|X_n - T_{xn}\|^2 \dots \quad (7)$$

$$\|Y_n - p\|^2 = (1-d_n) \|x_n - p\|^2 + d_n \|T_{xn} - p\|^2 - d_n (1-d_n) \|x_n - T_{xn}\|^2 \dots \quad (8)$$

Also by (2)

$$\|T_{yn} - p\| \leq k \max \{ \|y_n - p\|, \|y_n - T_{yn}\| \}$$

and

$$\|T_{xn} - p\| \leq k \max \{ \|x_n - p\|, \|x_n - T_{xn}\| \}$$

Let $S_1 = \{n \in \mathbb{N} : \|T_{yn} - p\| \leq k \|Y_n - p\|\}$

and $S_2 = \{n \in \mathbb{N} : \|T_{yn} - p\| \leq k \|Y_n - T_{yn}\|\}$

where N denote the set of positive integers.

Obviously $S_1 \cup S_2 = \mathbb{N}$

Suppose $n \in S_1$. Then using (8) we have

$$\begin{aligned} \|T_{yn} - p\|^2 &\leq k^2 \|Y_n - p\|^2 \\ &= k^2 (1-d_n) \|x_n - p\|^2 + k^2 d_n \|T_{xn} - p\|^2 - k^2 d_n (1-d_n) \|X_n - T_{xn}\|^2 \dots \quad (10^*) \end{aligned}$$

If in (10) $\|T_{xn} - p\| \leq k \|x_n - p\|$ holds. Then form 10*.

$$\begin{aligned} \|T_{yn} - p\|^2 &\leq [k^2(1-d_n) + K^4 d_n] \|x_n - p\|^2 - k^2 d_n (1-d_n) \|x_n - T_{xn}\|^2 \\ &\leq \|x_n - p\|^2 - k^2 d_n (1-d_n) \|x_n - T_{xn}\|^2 \end{aligned}$$

Thus for all $n \in S_1$

$$k^2 c_n d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2 - \|x_{n+1} - p\|^2 \dots \quad (11)$$

Now supposing that $n \in S_2$ we have by (7)

$$\begin{aligned} \|T_{yn} - p\|^2 &\leq k^2 \|y_n - T_{yn}\|^2 \\ &= k^2 (1-d_n) \|X_n - T_{yn}\|^2 + k^2 d_n \|T_{xn} - T_{yn}\|^2 - k^2 d_n (1-d_n) \|x_n - T_{xn}\|^2 \dots \quad (12) \end{aligned}$$

Since $\|x_n - T_{yn}\| = d_n \|x_n - T_{xn}\|$

$$\|y_n - T_{yn}\| = (1-d_n) \|x_n - T_{xn}\| \text{ and}$$

T satisfies (*) we have

$$\|T_{xn} - T_{yn}\| \leq k \max \{ \|x_n - T_{xn}\|, \|y_n - T_{yn}\|, \|x_n - T_{xn}\| \} = k A_n$$

Where A_n denotes the maximum of the set.

$$\text{Let } S_2^1 = \{n \in S_2 : A_n = \|x_n - T_{xn}\|\}$$

$$S_2^{11} = \{n \in S_2 : A_n = \|y_n - T_{yn}\|\}$$

$$S_2^{111} = \{n \in S_2 : A_n = \|x_n - T_{yn}\|\}$$

Clearly $S_2 = S_2^1 \cup S_2^{11} \cup S_2^{111}$

Now if $n \in S_2^1$ i.e. if

$$\|T_{xn} - T_{yn}\| \leq k \|x_n - T_{xn}\|, \text{ then form } \quad (12)$$

$$\|T_{yn} - p\|^2 \leq k^2 (1-d_n) \|x_n - T_{yn}\|^2 - k^2 d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2$$

and using (6) we get

$$\begin{aligned} \|x_{n+1} - p\|^2 &\leq (1-c_n) \|x_n - p\|^2 + c_n k^2 (1-d_n) \|x_n - T_{yn}\|^2 - c_n k^2 d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2 \\ &= (1-c_n) \|x_n - p\|^2 - k^2 c_n d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2 - c_n [1-c_n - k^2 (1-d_n)] \|x_n - T_{yn}\|^2 \\ &\leq \|x_n - p\|^2 - k^2 c_n d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2 \end{aligned}$$

Since $k^2 (1-d_n) < (1-d_n) \leq (1-c_n)$

$$\text{or } k^2 c_n d_n (1-d_n - k^2) \|x_n - T_{xn}\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \quad (13)$$

If $n \in S_2^{11}$ i.e. if $\|T_{xn} - T_{yn}\| \leq k \|y_n - T_{yn}\|$

Then using (7) we obtain

$$\begin{aligned} \|T_{xn} - T_{yn}\|^2 &\leq k^2 \|y_n - T_{yn}\|^2 \\ &\leq k^2 (1-d_n) \|x_n - T_{yn}\|^2 + k^2 d_n \|T_{xn} - T_{yn}\|^2 \\ &\quad - k^2 d_n (1-d_n) \|x_n - T_{xn}\|^2 \end{aligned}$$

and hence

$$\begin{aligned} \|T_{xn} - T_{yn}\|^2 &\leq [k^2 (1-d_n) / (1-k^2 d_n)] \|x_n - T_{yn}\|^2 \\ &\quad - [k^2 d_n (1-d_n) / (1-k^2 d_n)] \|x_n - T_{yn}\|^2 \end{aligned}$$

Then by (12)

$$\|T_{y_n} - p\|^2 \leq k^2(1-d_n) [(1+k^2d_n / (1-k^2d_n)) \|x_n - T_{y_n}\|^2 - k^2d_n(1-d_n) [1+k^2d_n (1-k^2d_n)] \|x_n - T_{x_n}\|^2]$$

$$= [k^2(1-d_n) / (1-k^2d_n)] \|x_n - T_{y_n}\|^2 - [k^2d_n(1-d_n) / (1-k^2d_n)] \|x_n - T_{x_n}\|^2$$

From which we obtain on using (6)

$$\|x_{n+1} - p\|^2 \leq (1-c_n) \|x_n - p\|^2 + C_n [k^2 (1-d_n) / (1-k^2d_n)]$$

$$\|x_n - T_{y_n}\|^2 - c_n[k^2d_n (1-d_n) / (1-k^2d_n)] \|x_n - T_{x_n}\|^2 - c_n (1-c_n) \|x_n - T_{y_n}\|^2$$

$$= (1-c_n) \|x_n - p\|^2 - [k^2c_nd_n (1-d_n) / (1-k^2d_n)] \|x_n - T_{x_n}\|^2$$

$$- [c_n (1 - k^2 - c_n (1-k^2d_n)) / (1-k^2d_n)] \|x_n - T_{y_n}\|^2$$

Since $C_n \rightarrow 0$ as $n \rightarrow \infty$ there exists an $n_0 \in \mathbb{N}$ such that $1 - k^2 > c_n$ for all $n \geq n_0$. Thus $n \geq n_0$ the last form on the right hand sides of the above expression is positive and hence we get $n > n_0$.

$$[k^2c_nd_n (1-d_n) / (1-k^2d_n)] \|x_n - T_{x_n}\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2 \dots \quad (14)$$

If $n \in S_2''$ i.e. iff

$$\|T_{x_n} - T_{y_n}\| \leq k \|x_n - T_{y_n}\|, \text{ then form (6) and 12 we obtain.}$$

$$\|x_{n+1} - p\|^2 \leq (1-c_n) \|x_n - p\|^2 + C_n [k^2 (1-d_n) \|x_n - T_{y_n}\|^2 + k^2d_n k^2 \|x_n - T_{y_n}\|^2 - k^2d_n (1-d_n) \|x_n - T_{x_n}\|^2]$$

$$- c_n(1-c_n) \|x_n - T_{y_n}\|^2$$

Hence for $n \geq n_0$ we get as before

$$k^2c_nd_n (1-d_n) \|x_n - T_{x_n}\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2$$

Since $1-d_n-k^2 \leq 1-d_n \leq (1-d_n) / (1-k^2d_n)$ for all n .

We obtain from the inequalities (13) (14) and (15) that for all $n \in S_2$ and $n \geq n_0$

$$k^2c_nd_n (1-d_n-k^2) \|x_n - T_{x_n}\|^2 \leq \|x_n - p\|^2 - \|x_{n+1} - p\|^2$$

Since this inequality holds for all $n \in S_1$ sec (11) it follows that (11) holds for all $n \in \mathbb{N}, n \geq n_0$

Now choosing $m \geq n_0$ and adding the in equalities (11) for values $m, m + 1, \dots, n$ of n .

We obtain

$$\sum_{j=m}^n l^2 c_j d_j (1-d_j-k)^2 \|x_j - T_{x_j}\|^2 \leq \|x_m - p\|^2 - \|x_{n+1} - p\|^2 \dots \quad (16)$$

Since $d_j \rightarrow 0$ as $J \rightarrow \infty$, $k^2 (1 - d_j - k^2)$ is +ve and bounded away from zero. The fact that right hand side of the above inequality is bounded and that $\sum_{j=1}^{\infty} c_j d_j = \infty$

imply that $\lim_{n \rightarrow \infty} \|X_n - T_{x_n}\| = 0$. Hence by compactness of E . It follows that there exists a subsequence (X_{n_k}) such that $\lim_{n \rightarrow \infty} X_{n_k} = q$ and $\|q - T_q\| = 0$ i.e. $T_q = q$. But by uniqueness of a fixed point of T , $q = p$. Again since the sequence $(\|x_n - p\|)$ is monotonically decreasing (as evident from (11)) and $\lim_{n \rightarrow \infty} X_{n_k} = p$ it ultimately follows that $\lim_{n \rightarrow \infty} X_n = p$.

This completes the proof.

1.3 FURTHER GENERALISATION

B. Fisher [7p, 8p] established the existence of a common fixed point of a pair of commuting mapping S and T satisfying the inequality.

$$\|S_x - T_y\| \leq k \max \{ \|x - y\|, \|x - T_y\| \|y - S_x\| \|x - S_x\| \|y - T_y\| \}$$

She proved the following theorem.

Theorem - 1

Let S and T be commuting mapping of a complete metric space (X, d) into itself satisfying (2) (with $d(x, y) = \|x - y\|$ of course) for all X, Y in X , where $0 \leq k < 1$ and the inequality. $\sup \{d(S^{r+1}T^n x, S^r T^{n+1} X, S^r T^n x): r, n = 0, 1, 2, \dots\} < k$ For some particular X in X . Then S and T have a unique common fixed point Z . Further Z is the unique fixed point of S and T .

We know that from (16) an iteration involving two mapping S and T which satisfy (17) converges to their common fixed point. This result reduces to Theorem-1 in the case S and T .

Theorem - 2

Let E be a compact and convex subset of a Hilbert space H . Let S and T be a pair of commuting self mapping S on E satisfying the inequality (9) for all $x, y \in E$ and $0 \leq k < 1$. Let the sequence (X_n) be defined on E by the iteration $X_1 \in E, X_{n+1} = (1-c_n)X_n + C_n T[(1-d_n)X_n + d_n S X_n], n \geq 1, \dots, (18)$. Where (C_n) and (d_n) are real sequences satisfying conditions (1), (11) of theorem 1. Then (X_n) convergence to the common fixed point of S and T .

Proof

Since E is compact and S and T are commuting mappings satisfying (17), the conditions of Theorem A of Fisher are satisfied where there exists a unique common fixed point P . Say of S and T . The proof of convergence of (X_n) to P is similar to the proof of Theorem 1 and hence omitted.

1.4 CONCLUSION

Iteration (18) of theorem 2 above shows that Ishikawa iteration scheme I (X_1, C_n, D_n, T) can be generalized by introducing more number of mapping which can be used to yield common fixed point (s) of the mappings Investigation in this direction has been carried out in Chapter III and Chapter IV.

REFERENCES

1. Banach, S. Sures operations dans les ensembles abstraits. *Learn applications aux equation integrals fund, Math*, (3) (1922).
2. Kirk, W.A.: Some recent results in metric fixed point theory *Journal of fixed point theory and application* (2007).
3. Hu, L-G: strong convergence of a modified Halpern's iteration for non expansive mappings. *Fixed point theory and application* (2008).
4. Bose, R.K and Mukherjee, R.N. Stability of fixed points sets and common fixed points of families of mapping *Indian J. Pure. Appl. Math* 11(9) (1980).
5. Bose, R.K and Mukherjee, R.N. Approximating fixed points of some mapping. *Proc. Amer. Math.Soc.* 82(4) (1981) 603-606.
6. M. Elin, M. Levenshtein, S. Reich and D. Shoikhet two rigidity theorems for holomorphic generators of continuous semi groups, *J. Non linear convex, Anal.* 9 (2008) (59-64).
7. Ceng, IC, Yao, JC: Hybrid viscosity approximation schemes for equilibrium problems and fixed point problems of infinitely many non expansive mappings. *Appl. Math. Comput.* 198, 729-741 (2008).
8. Hussain, N, Khamsi, MA: On asymptotic pointwise contractions in metric spaces. *Nonlinear Analysis: Theory, Methods and Applications.* 71 (10), 4423-4429 (2009).
9. S. Jain and V. H. Badshah, "Fixed Point Theorem of Multi-Valued Mappings in Cone Metric Spaces", *International Journal of Mathematical Archive*, Vol. 2, No. 12, 2011, pp. 2753-2756.
10. Reich, S, Sabach, S: Three strong convergence theorems regarding iterative methods for solving equilibrium problems in reflexive Banach spaces. *Optimization Theory and Related Topics.* 225-240 (2012).
11. N. Chen and J. Q. Chen, "Operator Equation and Application of Variational Iterative Method", *Applied Mathematics*, Vol. 3, No. 8, 2012, pp. 857-863. Doi:10.4236/am.2012.38127.
12. Chen, M. Pand Shin, M.H. Fixed points theorems for point to point and point to

- set maps, J. Math. Anal. Appl. 71 (1979) 516-524.
13. Ćirić, L.B. A generalization of Banach contraction principle. Proc. Amer-Math. Soc. 45(2) 1974 (267-273).
14. Ćirić, L, quasi contractions in Banach's spaces publ. del' inst. Math. 21 (35) (1977) 41-48.
15. S. Aizicovici, S. Reich and A.J. Zaslavski, Minimizing convex functions by continuous descent methods, Electron. J. Differ. Equ. 19 (2010), 7 pp.
16. Wang, S: A note on strong convergence of a modified Halpern's iteration for nonexpansive mappings. Fixed Point Theory and Applications, 2010 (2010).
17. Rhoades, B.E., A comparison of various definitions of contractive mappings. Trans, Amer, math. soc., 266 (1977) 257-290.

How to cite this article: Mishra K, Pattnaik H. Fixed points of quasi contractive mapping using ishikawa iteration. Int J Res Rev. 2016; 3(7):45-49.

